


Overall complications (RCT)


RCT: DTI vs. TE implant-based breast reconstruction, overall complications



*Adjusted OR: 3.42 (95% CI: 2.28-5.12)


Overall complications (NRSIs)

NRSI: DTI vs. TE implant-based breast reconstruction, overall complications



Reoperation (RCT)


RCT: DTI vs. TE implant-based breast reconstruction, reoperation



*Adjusted OR: 2.91 (95% CI: 1.34- 6.34)

Reoperation (NRSIs)

NRSI: DTI vs. TE implant-based breast reconstruction, reoperation




BREAST-Q dimensions, only measurements from 2 years onwards after reconstruction (according to median follow-up or time point of QoL measurement)


Notes: In Aliotta 2021, BREAST-Q scores were reported separately for tissue expander groups with and without ADM (read from graph). For the analysis in this report, groups were combined using this [resource / formula by Cochrane](#) (Practical 2 on how to include multi-arm studies in meta-analysis) by Emma Axon.

Alle Beobachtungszeiträume für die BREAST-Q Skalen liegen zwischen medianen/durchschnittlichen 6 Monaten und 36 Monaten; häufig mit einem längeren Abstand der Messung zur Operation für die DTI-Gruppe.


Satisfaction with breast (RCT)



Satisfaction with breast (NRSIs)



Satisfaction with overall outcome (RCT)



Satisfaction with overall outcome (NRSIs)

